

# RUSSELL M. LAFRANCE MD

Orthopedic Surgeon & Sports Medicine

### ACTIVITY:

- You should flex and extend your wrist and elbow several times throughout the day.
- Wear the sling for comfort, it is ok to come out of the sling and start gently moving your shoulder.
- Do not lift heavy objects with your operative arm until you are seen in follow up.
- Physical Therapy can start next week

### DIET:

- Start with liquids and advance your diet as tolerated.

### Medication:

- Take the prescription pain medicine (Norco/Vicodin/Percocet) as prescribed. DO NOT take Tylenol in addition to this.
- Ibuprofen or Advil can be taken in addition to the narcotic
- If you are taking prescription pain medication, you should not drive, operate machine/power tools, or drink any alcoholic beverages.

### DRESSING:

- Keep your dressing clean and dry. Leave the dressing intact until you see Dr. LaFrance in the office.
- Sponge baths only
- Use ice over your shoulder for 20 minutes intervals (no longer) for 3-4 times per day during the first week after your surgery.

### SPECIAL INSTRUCTIONS:

- Once you are off pain medications and have full strength and motion of your elbow, then you may drive.

### YOU SHOULD CALL YOUR DOCTOR FOR ANY OF THE FOLLOWING:

- Fever greater than 101.5° F
- Redness, warmth in the leg, knee, or calf
- Foul smelling drainage from the incision

## Clavicle Fracture ORIF – Patient Information

---

- Pain uncontrolled with pain medication as prescribed by your doctor
- Persistent nausea or vomiting into the next day
- Increased swelling of your fingers or severe tightness of bandage
- Chest pain or difficulty breathing

### **FOLLOW-UP CARE:**

- Your postoperative appointment 7-10 days after surgery will have been made for you when surgery was set up.